arXiv:0712.1484v1 [astro-ph] 10 Dec 2007

Frequency Dependence of Radio Images of Supernova Remnants

A.V. Karnaushenko[1], *E. Yu. Bannikova*[1,2], *V. M. Kontorovich*[1,2]

[1] Institute of Radio Astonomy NAS of Ukraine
[2] Karazin Kharkiv National University
a_karnaushenko@mail.ru, bannikova@astron.kharkov.ua, vkont@ira.kharkov.ua

Radio images of supernova remnants in the framework of diffusion model are discussed. The distribution profiles of synchrotron radiation intensity for spherical injection source of relativistic electrons are reduced at different frequencies. An explanation of the observational data obtained on UTR-2, according to which the size of the supernova remnant at decametric waves is larger than the remnant size at high frequencies, is given.

Introduction

The radio images of the supernova remnants (SNR) obtained with the UTR-2 radio telescope by M.A. Sidorchuk and E.A. Abramenkov show the difference between SNR sizes at low frequencies (decametric waves) vs. the higher ones [1].

For example, the brightness temperature contour map of SNR HB3 at the frequency 20 MHz is shown in Fig.1. The temperatures on the isophotes are given in $10^3 K$. At the lower corner the UTR-2 directional diagram is shown as a dashed circle. The center of SNR is denoted by the white "+". At the frequency 1420 MHz radio image of SNR corresponds to the size 60×80 pc (the region limited by a white circumference)[2]. At frequency 20 MHz it corresponds to the size 70×95 pc [1]. We can see that the SNR size at low frequency is larger than the SNR size at high frequency. In this work we propose an explanation of the SNR size changing with frequency in the framework of diffusion model.

Diffusion model

In the diffusion model we consider the SNR as a spherical source of relativistic electrons. The particles are accelerated by the spherical shock wave (SW) front and next further propagate by means of diffusion into the environment medium losing their energy due to synchrotron radiation [3]. The basic equation is the kinetic equation (KE) for electron distribution function (EDF) $N(E, t, \vec{r})$ with a source $Q(E, t, \vec{r})$ and with diffusion as a mode of the electron propagation

$$\frac{\partial N}{\partial t} + \frac{\partial (B(E)N)}{\partial E} - D\Delta N = Q(E, t, \vec{r}), \tag{1}$$

were Δ is the Laplace operator, D is the diffusion factor that as we will show bellow can be chosen to have no energy dependence.

The second term in KE describes the synchrotron losses of electron

$$B(E) = -\beta E^2, \quad \beta = \frac{32\pi}{9}\left(\frac{e^2}{mc^2}\right)^2 \cdot \frac{W_H}{m^2c^3},$$

where e is the electron charge, m is the electron mass, c is the light velocity and $W_H = H^2/(8\pi)$ denotes the energy density of the magnetic field.

The right part of equation (1) corresponds to the source of relativistic electrons:

$$Q(E, t, \vec{r}) = Q_0 \cdot N_{inj}(E) \cdot S(\vec{r}) \cdot \Theta(t),$$

where $S(\vec{r}) = \delta(|\vec{r}| - R)$ sets the spherical form of the source, with radius R, $N_{inj} = E^{-\gamma_0} \cdot \Theta(E_{max} - E) \cdot \Theta(E - E_{min})$ is the distribution function of injected electrons, $\gamma_0 = 2$, $\Theta(...)$ is the Havyside's step function. The solution of KE is the EDF for the spherical source [3], [4]:

$$N(E,t,\vec{r}) = \frac{Q_0}{(4\pi)^{3/2} \cdot E^2} \cdot \int_{-\tau_+(E,t)}^{\tau_-(E,t)} d\tau'_- \cdot \frac{\tilde{E}^2(\tau_+, \tau'_-)}{\alpha^{3/2}(\tau_+, \tau'_-)} \cdot N_{inj}(\tilde{E}) \cdot R^2 \int d\Omega \cdot exp\left(\frac{(\vec{r} - \vec{r'})^2}{4 \cdot \alpha(\tau_+, \tau'_-)}\right), \tag{2}$$

where

$$\tau_\pm(E,t) = \frac{1}{2} \cdot \left(t \pm \frac{1}{\beta}\left(\frac{1}{E_{max}} - \frac{1}{E}\right)\right), \quad \alpha(\tau_+, \tau'_-) = D_0 \cdot (\tau_+ + \tau'_-),$$

$$\tilde{E}(\tau_+, \tau'_-) = \frac{1}{\frac{1}{E_{max}} - \beta \cdot (\tau_+ - \tau'_-)},$$

$(\vec{r} - \vec{r'})^2 = (x - x')^2 + (y - y')^2 + (z - z')^2$, $x' = R \cdot \sin\theta \cdot \cos\varphi$, $y' = R \cdot \sin\theta \sin\varphi$, $z' = R \cdot \cos\theta$, $d\Omega = \sin\theta d\theta d\varphi$.

The physical parameters of the diffusion model are the following: he life time $\tau = 1/(\beta \cdot E)$ - the time of particle existence with energy E, diffusion length $l_{dif} = \sqrt{D_0 \cdot \tau}$ - the distance, which particle passes during the life time, and the diffusion velocity is $V_{dif} = D_0/l_{dif}$. The diffusion length defines the size of the radio image of SNR. Knowing the EDF, we can calculate the intensity distribution of the synchrotron radiation I [5]. The intensity distribution in the image plane for the sources resolved by the radio telescopes is given by:

$$I(\nu, t, \vec{r}) = \frac{\sqrt{3} \cdot e^2}{m \cdot c^3} \int_0^{E_{max}} dE \int dz \cdot N(E,t,\vec{r}) \cdot H_\perp \cdot \frac{\nu}{\nu_s} \cdot \int_{\nu/\nu_s}^{\infty} K_{5/3}(\eta) d\eta, \tag{3}$$

$$\nu_s = \frac{3}{4\pi} \cdot \frac{eH}{mc} \cdot \left(\frac{E}{mc^2}\right)^2,$$

where $H_\perp$ is the magnetic field projection on the image plane, $\int dz$ is an integral along the line of sight and $K_{5/3}$ is modified Bessel function. The numerical calculations were made with help of the Mathematica 5.1 package.

Frequency dependence of synchrotron radiation intensity distribution

The profiles of synchrotron radiation intensity distribution at different frequencies are given in Fig.2. Though the remnant is not fully symmetrical we choose a sphere with $R = 50pc$ as approach. We neglect the motion of the shock front and consider the SNR as an immovable injection source.

In Fig.2a $\tau = 3.6 \cdot 10^{11}s, l_{dif} = 1.3 \cdot 10^{20}cm, V_{dif} = 3.7 \cdot 10^8 cm/s$: the diffusion length is higher than SNR radius, it almost completely defines the size of the radio image. In Fig.2b $\tau = 8 \cdot 10^{10}s, l_{dif} = 6.3 \cdot 10^{19}cm, V_{dif} = 7.8 \cdot 10^8 cm/s$, the life time and the diffusion length decrease, and a gap (valley) appears in the figure. In Fig.2c the life time and the diffusion length further decrease $\tau = 5 \cdot 10^{10}s, l_{dif} = 5 \cdot 10^{19}cm, V_{dif} = 9 \cdot 10^8 cm/s$, and the gap increases. In Fig.2d with parameters $\tau = 4.2 \cdot 10^{10}s, l_{dif} = 4.6 \cdot 10^{19}cm, V_{dif} = 1 \cdot 10^9 cm/s$ the gap reaches maximum value: the diffusion length becomes less than own size of the system and has no influence in its radio image. The approximation of a constant diffusion factor is valid if the following equality is carried out: $l_{dif1}/l_{dif2} \approx \sqrt[4]{\nu_2/\nu_1}$. For the chosen frequencies and corresponding values of diffusion lengths ($l_{dif1} = 1.3 \cdot 10^{20}cm, \nu_1 = 20MHz$ and $l_{dif2} = 4.6 \cdot 10^{19}cm, \nu_2 = 1420MHz$) the equality is carried out with a good accuracy $1.3 \cdot 10^{20}cm/4.6 \cdot 10^{19}cm \approx \sqrt[4]{1420MHz/20MHz} \approx 2.9$ It is necessary to note, that velocity of the SNR shock front $R(t) = 3 \cdot 10^7 cm/s$ can be of the same order as the diffusion velocity and even more, therefore in general it is necessary to consider the front movement.

Conclusions

In the framework of the diffusion model we have explained the observational data obtained by the UTR-2 according to which the size of the supernova remnants on the low frequencies is is higher than the one at high frequencies. It has been shown that in our case the diffusion factor may be considered as a constant.

Acknowledgements

The authors thank M.A. Sidorchuk for the image of SNR HB3.

References

[1] Abramenkov, E. A., Sidorchuk, M. A. Radiophysics and Radioastronomy, V. 11, 2, pp. 134-154 (2006)

[2] Tian, W. W., Leahy, D. A. Astronomy and Astrophysics, V. 436, pp. 187-193 (2005)

[3] Bannikova, E.Yu., Kontorovich, V.M., Radiophysics and Radioastronomy, V. 9, 1, pp. 29-36 (2004)

[4] Bannikova, E.Yu., The journal of Kharkiv National University, physical series. Nuclei, Particles, Fields, V. 627, pp. 57-62 (2005)

[5] Ginzburg, V. L., "Theoretical Physics and Astrophysics", Nauka, Moscow, 488 p. (1987)

Figure 1. Brightness temperature contour map of SNR HB3 at frequency 20 MHz by M.A. Sidorchuk and E.A. Abramenkov [1].

Figure 2. Profiles of intensity of synchrotron radiation distribution for spherical source with constant diffusion factor for following parameters a) $\nu = 20MHz$; b) $\nu = 408MHz$; c) $\nu = 1000MHz$; d) $\nu = 1420MHz$, I_ν is expressed in relative units in the same scale.

Figures are available on YSC home page (http://ysc.kiev.ua/abs/proc14_8.pdf).